



Review Article

REVIEW OF LANGALI (*GLORIOSA SUPERBA* LINN) – AGADA TANTRA PERSPECTIVE

Athulya.C.M^{1*}, Neethu P¹, Vijitha Vijayan¹, Arathi Rajesh²

*12nd Year PG Scholars, ²Associate Professor, Dept. of Agadatantra, MVR AMC, Parassinikadav, Kannur, Kerala, India.

ABSTRACT

Agadatantra is a branch of Ayurveda which deals with the symptoms and treatment of various animal, plant, artificial and latent poisons. The drug *Langali* (*Gloriosa superba* Linn) is being categorized under *Moolavisha* (poisonous roots) and is also one among the nine *Upavishas* (semi poisonous drugs) by different Ayurvedic pharmacopoeias. Even though it is poisonous it has got *Shothahara* (anti-inflammatory), *Vranahara* (wound healing), *Krimigna* (wormicidal), *Shalyaharana* (removal of foreign substances), *Garbapaatana* (abortifacient) properties etc. It is being used by the Ayurvedic physicians in treatment of various ailments after proper purification. But its purificatory methods are not mentioned in Samhitas (treatises). It is mentioned in Ayurveda prakasha and traditional Agadatantra text books of Kerala. Here in this article an attempt is made to collect the information regarding *Langali* in *Agadatantra* perspective.

The article includes both modern and Ayurvedic concepts in regards to its symptoms during poisoning and its treatment. Details regarding its purificatory methods and its therapeutic use in compound formulations as an antidote which has been mentioned in the traditional Agadatantra books which are available in Kerala have also been included.

KEYWORDS: Agadatantra, Ayurveda, *Langali*, *Upavisha*, *Moola Visha*.

INTRODUCTION

Agadatantra is a branch of Ayurveda which deals with the symptoms and treatment of various animal, plant, artificial, latent poisons. The drug *Langali* (*Gloriosa superba* Linn) has been categorized as one of the nine *Upavishas* (semipoisonous) of Ayurvedic pharmacopoeias. *Langali* is commonly used as an abortifacient. It is used in various compound formulations in the treatment of poisoning. In this article an attempt is made to collect the information regarding its poisonous effects, treatment and its therapeutic usage after purification.

Scientific Classification ^[1]

Kingdom	Plantae
Subkingdom	Tracheobionta
Division	Spermatophyta
Class	Liliopsida
Subclass	Liliidae
Order	Liliales
Family	Liliaceae
Genus	Gloriosa
Species	Gloriosasuperba Linn.

Botanical Description ^[1]

A perennial herbaceous climbing herb growing 3.5 to 6.0 meters in length.

Tubers

- V or L-shaped, finger-like that are pure white when young, becoming brown with age.

Leaves

- Pointed, dark green, glossy, each equipped with a tendril by means of which it clings onto other plants.
- Occur in whorls of 3 to 4, opposite or alternate, simple, sessile, ovate to lanceolate ranging from 6 to 20 cm in length and 1.5 to 4 cm wide.

Flowers

- Attractive, borne on long stalks.
- Have six erect petals ranging in colour from bright yellow to bicoloured, red and yellow or purple and yellow.
- The genus derives its name from the Latin word 'gloriosus', referring to the flowers.

Fruits

- Capsules that split open to release several smooth red seeds with a spongytest (Maroyi et al., 2011).

Vernacular Names [2]

English	Malabar glory lily
Hindi	<i>Karihari, Languly</i>
Kannada	<i>Kolikuttuma</i>
Malayalam	<i>Mentonni, Mettonni, Kantal</i>
Sanskrit	<i>Langali, Vishalya</i>
Tamil	<i>Kalappaikkilanku, Nabhikkodi</i>
Telugu	<i>Adavinabhi</i>

Habitat of *G. superba* Linn. [1]

Gloriosa is a native of tropical Asia and Africa. Tropical India: North -West Himalayas to Assam and the Deccan peninsula, extending up to an elevation of 2120 M.

South India: In Karnataka, commonly found growing all along the Western Ghats.

Others: Madagascar, Srilanka, Indo-China and on the adjacent island. (Faroogi and Sreeramu, 2001).

Chemical Constitutes of *G.superba* Linn. [1]

Studies reveal that all parts of the plant, especially the tubers are extremely toxic due to the presence of a highly active alkaloid, Colchicine. The species also contains another toxic alkaloid, Gloriosine. Other compounds which are isolated from the plants are lumicolchicine, 3-demethyl-N-deformyl-N-deacetylcolchicine, 3-demethyl colchicine, N-formyldeacetyl- colchicine (Chulabhorn et al., 1998). The tubers or dried roots contain benzoic and salicylic acid, sterols and resinous substances-colchicines, 3-demethyl colchicine, 1, 2-didemethyl colchicine, 2, 3- didemethyl colchicine, N-formyl, N-deacetylcolchicines, colchicocide, tannins and superbine (Jain and Suryavanshi, 2010). Flower's contain Luteolin, its Glucoside, N-Formyl-de-Me-colchicine, its Glucoside and 2-de-Me-colchicine Content of Colchicine 0.25%.

Metabolism of Colchicine [9]

- Colchicines are rapidly absorbed from the intestine and undergoes significant first -pass hepatic metabolism.
- Excretion is through faeces.
- Renal clearance accounts for about 10-20% of colchicines excretion.
- Colchicines inhibit the polymerization of microtubules and formation of mitotic spindle in cell division. Therefore, the rapidly dividing cells of the intestinal mucosa are severely affected.

Symptoms of poisoning [9]

- Colchicines can cause severe gastroenteritis, shock, multiple organ failure, electrolyte imbalance, metabolic acidosis, pancytopenia, hypotension, rhabdomyolysis, hypocalcaemia,

and adult respiratory distress syndrome, ascending polyneuropathy.

- Primary symptoms develop within two to six hours. Intense vomiting, numbness and tingling around the mouth, burning, and rawness of the throat, nausea, abdominal pain and bloody diarrhea leading to dehydration etc.
- Late manifestations that develop about one to two weeks after poisoning are alopecia and dermatitis.

Medico legal importance

Accidental:[31] Mostly, Due to its use in therapeutic medicine.

For criminal abortion it is introduced in vagina.

Homicidal, Suicidal: Rare

Fatal Dose: [21] 60 mg in adults

Fatal Period: [21] 12 to 72 hours

Treatment According to Modern [17]

- The patient requires immediate hospitalization, followed by gastric lavage.
- There is no specific antidote which is available for the treatment.
- Fluid resuscitation and/or ionotropic support for correction of fluid loss.
- Assisted ventilation for patients with respiratory depression.
- Dialysis for patients with renal failure and oliguria.
- Prophylactic antibiotic therapy is advisable if leucopaenia is present.
- Vitamin K and fresh frozen plasma should be given if the clotting time is abnormal.

Ayurvedic concept**Gana**

Nighantus (lexicons) and classical texts of Ayurveda details *Langali* under different categories by keeping the drug under various *Vargas* which are listed below.

Charakacharya mentioned it under Moolavisha (Group of root poisons) [3]

<i>Bhavaprakasha nigantu</i>	<i>Guduchyaadi varga</i> [4]
<i>Raaja nigantu</i>	<i>Shatahwadi varga</i> [5]
<i>Kaiyadeva nigantu</i>	<i>Oshadi varga</i> [6]
<i>Madanapala nigantu</i>	<i>Guduchyadi varga</i> [7]
<i>Dhanwantari nigantu</i>	<i>Karaveeradi varga</i> [8]

Ayurvedic texts which have classified *Langali* under *Upavishavarga* (Group of semipoisonous plants) are[30]

- Sharangadara samhita
- Bhava prakasha

- Yoga ratnakara
- Rasarnava
- Rasa ratnasamuchayam
- Rasa tarangini
- Rasa sanketikalika
- Rasendra chuudamani
- Ayurveda prakasha

Synonyms

Synonyms	B.P ^[4]	Ra.Ni ^[5]	K.Ni ^[6]	Ma.Pa ^[7]	Dha.Ni ^[8]
<i>Kalihari</i>	+	-	+	-	-
<i>Halini</i>	+	+	+	+	+
<i>Langali</i>	+	-	+	+	+
<i>Shakrapushpi</i>	+	-	-	-	-
<i>Vishalya</i>	+	+	+	+	+
<i>Agnishikha</i>	+	-	+	-	-
<i>Anantha</i>	+	-	-	-	-
<i>Vahnivaktra</i>	+	-	-	-	-
<i>Garbanut</i>	+	-	-	-	-
<i>Kalikaari</i>	-	+	-	+	+
<i>Langalini</i>	-	+	-	-	-
<i>Garbapaatini</i>	-	+	+	+	+
<i>Deepti</i>	-	+	-	-	-
<i>Agnimukhi</i>	-	+	-	-	+
<i>Hali</i>	-	+	-	-	-
<i>Nakta</i>	-	+	-	-	+
<i>Indupushpika</i>	-	+	-	-	+
<i>Vidyujjala</i>	-	+	-	-	-
<i>Agnijihwa</i>	-	+	-	+	-
<i>Vranahrut</i>	-	+	-	-	-
<i>Pushpasaurabha</i>	-	+	-	-	-
<i>Swarnapushpa</i>	-	+	-	-	-
<i>Vahnishikha</i>	-	+	-	+	-
<i>Seeri</i>	-	-	+	+	+
<i>Indrapushpi</i>	-	-	+	-	-
<i>Vahnijihwa</i>	-	-	+	-	-
<i>Pradepta</i>	-	-	+	-	-
<i>Shikha</i>	-	-	+	-	-
<i>Vahnimukhi</i>	-	-	+	+	-
<i>Prabhata</i>	-	-	+	+	-
<i>Pushpaseekara</i>	-	-	+	-	-
<i>Shuklapushpika</i>	-	-	-	+	-
<i>Vidyud</i>	-	-	-	+	-
<i>Ulka</i>	-	-	-	+	-
<i>Pushpasi</i>	-	-	-	+	-
<i>Bhara</i>	-	-	-	+	-
<i>Agnika</i>	-	-	-	+	-
<i>Nalarandri</i>	-	-	-	+	-
<i>Deepta</i>	-	-	-	-	+

Properties

Properties	B.p [4]	Ra.ni [5]	K.ni [6]	Ma.ni [7]	Dha.ni [8]
Rasa	Tikta Katu kashaya	Katu	Katu Tikta		Katu Tikta
Guna	Teekshna Laghu Kshara	-	Kshara Sara Teekshna Laghu	Sara Teekshna Laghu	Sara
Veerya	Ushna	Ushna	Ushna	Ushna	Ushna
Vipaka					
Doshakarma	Sleshmajit pittajanaka	Kaphavaatahara	Pittala	Pittala	Kaphavata vinashini
Rogagnata	Atisara Kushta Shopha Arsha Vrana Shola Krumi	Shalyanishkasakarini Saarini para	Vastishoola Balasakushta Shopha Arsha Vrana Jantuvinashini	Kushta Shopha Arsha Vrana Shoola Krumi	Shwayadhu Shalya Vrana
Causes	Garbapaata	Garbanta	Garbapaata	Garbapaatini	Garbapaata

Shodhana (Purification) of Ashudha (Toxic) Langali.^[12]

1. Soak Langali in Gomootra (Cow's urine) for 1 day or
2. Soak Langalikanda in Gomootra (cow's urine) for 1 day;
Then soak in Kwada (decoction) of Amruta (*Tinospora cordifolia*) and Tanduleeyakamoola (Root of *Amaranthus spinosus* L) for 30 nazika (30×24 minutes). Then in Godugda (cow's milk) for 1 day.

Symptoms of Intake of Langali Kanda Visha According to Ayurveda.^[29]

Chardi (vomiting), Kampa (tremors), can even lead to death.

Treatment For Langali Sevana according to Ayurveda (Kriya Koumudi)^[10]

1. Intake of Chanampayar (*Lens culinaris*), Bhumyamalaki (*Phyllanthus niruri*), Ardraka (ginger) in equal quantity.
2. Ardrakasevana (consumption of ginger) alone cures Visha.
3. Intake of Kwatha (decoction) of Bibitakatwak (*Terminalia bellerica* bark).
4. Intake of Shigrumoolatwakswarasa (juice of *Moringa oleifera* root bark) and Dadhi (curd) in equal quantity.
5. Intake and external application of Neelimoola (root of *Indigofera tinctoria*) and Maricha (*Piper nigrum*).

6. Haritakikwatha sevana (intake of *Terminalia chebula* decoction).

7. Intake of Shigrutwak (bark of *Moringa oleifera*) with Madhu (honey).
8. Intake of Shuntikalka (paste of dry ginger) with Ushnajala (hot water) or Sarvangalepa (whole body application).

Treatment According to Prayoga Samuchayam and Vishavaidya Jyotsnika.^[11]

1. Neelimoola (root of *Indigofera tinctoria*) with Maricha (*Piper nigrum*).

Kodashery Margam^[29]- Intake of Shunti (dry ginger) in boiled water.

Use of Langali in Agada prayoga

1. Saarkarmikaagada - Karaveera, Arkakusuma, Langalimoola, Kakana, Pata, Marichamade into Kalka (paste) with Aranala - for Vyantaradashta (Vyantara type of snake).^[13]
2. To prevent spread of Visha- Langalikanda, Madhuka, Ashwaganda, Nimbatwak, Draksha, Karam, Hingu, Chitraka, Shunti, Haridra, Mayakku, Chandana, Devadaruchurna triturated in Jambeera swarasa is made into Gutika (pills) and used for external application.^[14]
3. Peetanetramandali (variety of viper) Kattuchanakam, Langali, Kakanasa.^[15]
4. Raga mandali (viper variety)- Langali, Shireesha pushpa, moola - external application.^[16]
5. Swedana of Visharta (to induce sweating of poison inflicted person).

- a) *Langalipatra* alone or with combination of any drugs like *Erandapatra*, *Nimbapatra*, *Karanga patra* or *Shireeshapatra* boil in *Tandulodaka* and reduce to 1/4th and is used for *Swedana*.^[18]
- b) *Bakula*, *Langali*, *Musali*, *Shireesha*, *Nimbapatra* in *Tandulodaka*.^[25]
6. For all *Sarpavisha* (snake bites)– Root of *Langali* with *Maricha* (*Piper nigrum*) in equal quantity with pure water.^[22]
7. For *Nasya* (nasal instillation) and *Lepa* (external application) to cure *Moha* (unconsciousness) and *Visha* (poison)- *Kupilumajja* resin is extracted and kept in sunlight for 7 days; *Vacha*, *Hingu*, *Langalikanda* is taken in half quantity and *Mardana* (trituated) for 3 days; ¼th *Somnadihingu* is added and trituated with *Snuhiksheera* for 3 days and store in coconut shell. ^[23]
8. *Vishamoha* (to cure confused state due to *Visha*) - *Langalimoola* with *Maricha* made into *Gutika* (pills) and intake in pure water.^[24]
9. *Visha*, *Vatavyadinashana*- Intake of *Ankola*, *Eranda*, *Langali*, *Ishwaramooli*, *Nimba*, *Snuhi*, *Neeli*, *Shirisha*, *Arka*, *Dhatura*, *Bakula patras*, *Musali*, *Mundi* with *Tandulodaka* boiled and reduced to 1/4th.^[26]
10. *Mooshikavishakarnikapatana* (growth like pericarp of lotus in rat poisoning) - External application of *Langali*, *Alabumoola*, *Trivrut*, *Snuhi*, *Neelimoola*, *Apamarga* made into *Kalka* with *Tila*.^[27]

Neutralization effect against the venoms of *Naja nigricollis* venom

Evaluated the antivenom properties displayed by the alcoholic extracts of *Andrographis paniculata*, *Crateva magna*, *Gloriosa superba* and *Hydrocotyle javanica*. Extracts of *H. javanica* and *G. superba* gave 80 % and 90 % protection to mice treated with minimum lethal dose of venom (LD99). These two plants showed significant neutralization effect against the venoms of *Naja nigricollis* venom. *H. javanica* and *G. superba* (25-100 mg/mL) produced significant changes of membrane stabilization of human red blood cells (HRBC) exposed to hyposaline-induced haemolysis.^[19]

Antioxidant Activity and Anticancer Study against Human Cancer Cell (Hep-G2)

The *G. superba* methanolic extracts by In vitro Assay for Cytotoxicity Activity (MTT Assay) reveals the presence of different types of phyto constituents which has the capacity of anti-oxidant and cytotoxicity effect on Hep-G2 cells. Thus *G. superba* has the potentiality to inhibit the human carcinoma cell line growth.^[20]

Inhibition of the toxic effect of rattle snake venom

Traditional approach was evaluated scientifically with some selected plant extracts (7.2 mg/kg bw) and partially purified fractions (2.4 mg/kg bw) were orally administered to mice experimentally envenomed with rattlesnake venom S.C. injection (2.5–15µg/kg bw). The purified fractions (2.4mg/kg bw) produced significant protection against venom induced changes in serum SOD and LPx levels. The isolated fractions effectively inhibited the toxic effect of snake venoms in vitro than in vivo. The above observations confirmed the protective activity of plants- *Aristolochia indica*, *Hemidesmus indicus*, *Gloriosa superba*, *Strychnos nuxvomica*, *Eclipta prostrata*, and *Andrographis paniculata* against the lethal action of snake venom.^[28]

Shodana (purification) of *Langali*

Purificatory measures are mentioned in other *Grandas* in which *Takra* and *Saindhava* are made into solution and pieces of *Langali* are immersed in it for 3 days or 7 days.^[33]

UV spectroscopy method was used to analyze the media (*Takra* and *Saindhava*) before and after *Shodhana* to provide objectiveness to the process of *Shodhana* and role of media in the process. Loss on drying, total ash, water soluble extractive and alcohol soluble extractive were found to be increased in the samples after *Shodhana* with the media as compared to the samples before *Shodhana*, while acid insoluble ash was found to be decreased after *Shodhana* procedure. Qualitative analysis shows presence of alkaloids, saponins, tannins and carbohydrates in the samples before and after *Shodhana* although their quantification was not done. The article suggest further studies in the purification using different media.^[34]

Abortifacient property and Cardiovascular effects

Phytochemical analysis as well as toxicity (in mice) and antifertility studies, uterotrophic assay, duciduomamodel, uterotonic assessment in-vitro and in-vivo of aqueous extract was carried out in rats. Oxytocin was used as the uterotonic reference standard. The effect of the extract on cardiovascular parameters was also evaluated. Phytochemical analysis shows presence of flavonoids, tannins, alkaloids, and glycosides in the aqueous extract of *Gloriosa superba*. The aqueous extract of *Gloriosa superba* showed oxytocic activity and early abortifacient activity which may be due to the presence of alkaloids such as colchicine. Both the reference (oxytocin) and the extract produced dose dependent contractions but the extract had no effect on heart parameters and blood pressure.^[35]

DISCUSSION

Even an acute poison can become an excellent drug if it is properly administered. On the other hand even a drug if not properly administered, becomes an acute poison.^[32] This same principle applies in case of *Langali* also. Properly used *Langali* acts as medicine, while improper use may even lead to death. It is being categorized under *Moolavisha* (poisonous roots) and one among the nine *Upavishas* (semi poisonous drugs) by different Ayurvedic pharmacopoeias. *Langali* is used in Ayurveda for curing many diseases. *Langali* is having *Katu, Tikta rasa* (pungent, bitter); *Ksharaguna* (alkaline property), *Ushnaveerya* (hot in potency), *Teekshna* (sharp), *Laghu* (lightness), *Sara* (moving) *Guna* (property). It is having *Vatakaphashamana property* (pacifies *Vatakaphadosha*) and is *Pittavardaka* (increases *Pitta*). It is used for curing *Shotha* (inflammation), *Vrana* (wounds), *Krumi* (anti-microbial), *Kushta* (skin disorders) etc. *Keraleeyavisha Vaidya granthas* (toxicological texts) have given the details of the purificatory measures to be adopted for *Langali* and treatment that has to be given in case of its poisoning.

CONCLUSION

Researchers have shown the antivenom potential of *gloriosa* against cobra venom and also its anti-oxidant property. It is the need for the hour to carry out further researches regarding the toxic and anti-toxic properties of the drug.

Acknowledgement

Department of Agadatantra, MVR AMC, Parassinikadavu, Kannur, Kerala.

REFERENCE

1. Dharmendra singh et al. *Gloriosa superba* linn: An important endangered medicinal plant and their conservation strategies. International Journal of Botany and Research (IJBR). Mar 2013; Vol. 3 (1):19-26.
2. PK Varrier. Indian medicinal plants. Volume 3. Chennai; Orient longman private Ltd; Reprint 2007.p.76.
3. Ram Karan Sharma, Vaidya Bhagwan Dash. Carakasamhita of Agnivesa with Ayurveda dipika commentary by Cakrapani, Volume 4 (Chikitsa sthana). Varanasi; Chaukamba surbharati prakashan; 2017.p.325.
4. Sri.Brahmasankara Mishra, Sri.Rupalalajivisya. Bhavprakash. Varanasi; Chaukamba Sanskrit Sansthan; ninth edition 1999.p.313, sloka 80, 81.
5. Dr.Indradev Tripathi. Rajanighantu of Panditnarahari, edited with Dravyaguna prakashika Hindi commentary. Varanasi;

- Chaukambha Krishnadas Academy;p.86, sloka no 128-130.
6. Prof.Priyavratasarma, Dr.Guruprasada Sarma. Kaiyadeva Nighantu (pathyaapathyavibodhaka). Varanasi; Chaukambha Orientalia; first edition 1979.p.198-199, sloka no 1075-1077.
7. Ramaprasadpatiyala. Nrupamadanapalaviachita Madanapala Nigantu. Mumbai; Khemraj Srikrishadas Prakasan; November 2004.p.61, sloka no 28-30.
8. Acharya PV Sharma. Dhanwantari Nigantu. Varanasi; Chaukhambha Orientalia;p.122, sloka no 8, 9.
9. G. Prabhu et al. Clinical profile and outcome of gloriosa superba poisoning. Paripex - Indian journal of research. August 2016; 5 (8):1-3.
10. VM Kuttykrishnanmenon. Kriyakoumudi. Kottayam; Kerala sahityapravartaka co- operative society ltd;1986.p.752, sloka no 760- 764.
11. Kochunni Tamburan. Prayogasamucchaya (11th paricheda). Trissur; Salubha Books;1999.p.296. K. Mahadevashastri. Visha Vaidya Jyotsnika (13th adikara). Trivandrum; Authority of the government of his highness the maharaja of Travancore; p. 81, sloka no14.
12. VM Kuttykrishnanmenon. Kriyakoumudi. Kottayam; Kerala sahityapravartaka co- operative society Ltd;1986.p. 859.
13. Prof.K.R.Srikantamoorthy. Vagbata ashtanga hrudayam vol3 (uttarasthana). Varanasi; Choukamba krushnadas academy; 2015. p.354; sloka no 70, 71.
14. Dr.K.P.Sreekumari Amma. Kodasherymargam. Trivandrum; Trivandrum Publication division; 1988.p. 49.
15. Kochunni Tamburan. Prayogasamucchaya (3rd paricheda). Trissur; Salubha Books; 1999.p.66.
16. Kochunni Tamburan. Prayogasamucchaya (3rdparicheda). Trissur; Salubha Books; 1999.p.66.
17. Peter George MD. Death related to Herbal Therapy for Joint Pains – A rare case of *Gloriossa Superba* Poisoning. Journal of Clinical and Diagnostic Research. April 2011;5 (2):379 - 380.
18. K. Mahadevashastri. Visha Vaidya Jyotsnika (8th adikara). Trivandrum; Authority of the government of his highness the maharaja of Travancore; p. 46.
19. Chidambaram Kumarappan et al. Antihemolytic and snake venom neutralizing effect of some Indian medicinal plants. Asian Pacific Journal of Tropical Medicine. 2011;4 (9):743-747.
20. Simon SE et al. Antioxidant Activity and Anticancer Study on Phytochemicals Extract from Tubers of

- Gloriosa superba against Human Cancer Cell (Hep-G2). Journal of Pharmacognosy and Phytochemistry. October-December 2016;4 (4):7-12.
21. Peranantham S et al. Fatal gloriosa superba poisoning - A case report. International Journal of Medical and Pharmacological Sciences. June 2014; 04 (10):21-24.
22. Kuttikrishnanmenon. V.M.Kriyakaumudi. Kottayam; Sahitya Pravaraka Cooperative Society Ltd;1986.p.222;Sloka-172.
23. Kuttikrishnanmenon. V.M.Kriyakaumudi. Kottayam; Sahitya Pravaraka Cooperative Society Ltd;1986. p.340-341;Sloka-1727- 1730.
24. Kuttikrishnanmenon. V.M.Kriyakaumudi. Kottayam; Sahitya Pravaraka Cooperative Society Ltd; 1986.p.378 ;Sloka-382.
25. Kuttikrishnanmenon. V.M.Kriyakaumudi. Kottayam; Sahitya Pravaraka Cooperative Society Ltd; 1986. p.404;Sloka-124.
26. Kuttikrishnanmenon. V.M.Kriyakaumudi. Kottayam; Sahitya Pravaraka Cooperative Society Ltd; 1986. p.405;Sloka-127-129.
27. Kuttikrishnanmenon. V.M.Kriyakaumudi. Kottayam; Sahitya Pravaraka Cooperative Society Ltd;1986. p.506;Sloka-848-849
28. Samy et al. Ethnobotanical survey of folk plants for the treatment of snakebites in Southern part of Tamilnadu, India. Journal of Ethnopharmacology. 17 January 2008; 115 (2): 302-312.
29. Kodasherymargam, Edited by Dr.K.P.Sreekumari Amma; 1st edition; 1988; Publication division, Trivandrum; Page no 94.
30. Dr S.G.Huparikar, Dr V.P Joglekar; Text book of Agadatantra. Pune; Rashtriyashikshan mandal; 2008.p.20.
31. Dr S.G.Huparikar, Dr V.P Joglekar; Text book of Agadatantra. Pune; Rashtriyashikshan mandal; 2008.p.159.
32. Ram Karan Sharma, Vaidya Bhagwan Dash. Carakasamhita of Agnivesa with Ayurvedadipika commentary by Cakrapani, Volume 1 (Sootrasthana). Varanasi; Chaukamba surbharati prakashan; 2015.p.60.
33. Bhide Bhargav, Acharya Rabinarayan. Langali (Gloriosa superba) Linn. and its therapeutic importance in Ayurveda - a review. International Journal of Ayurvedic Medicine, 2012, 3 (2):58-67.
34. Bhide Bhargav, Shukla V J, Acharya Rabinarayan. UV Spectroscopic Analysis of Media (Takra and Saindhava) Used for Sodhana of Langali (Gloriosa superba Linn.) Root. Inventi Impact: Planta Activa, 2013 (1): 30-33.
35. Arati A Malpani et al. Effect of the Aqueous Extract of Gloriosa superb Linn (Langli) Roots on Reproductive System and Cardiovascular Parameters in Female Rats. Tropical Journal of Pharmaceutical Research; April 2011; 10 (2): 169-176.

Cite this article as:

Athulya.C.M, Neethu P, Vijitha Vijayan, Arathi Rajesh. Review of Langali (Gloriosa Superba Linn) - Agada Tantra Perspective. International Journal of Ayurveda and Pharma Research. 2018;6(11):51-57.

Source of support: Nil, Conflict of interest: None Declared

***Address for correspondence
Dr.Athulya. C.M**

2nd Year PG Scholars,
Dept. of Agadatantra, MVR AMC,
Parassinikadav, Kannur, Kerala,
India.
Cell: 9400307721
Email: cmathulya@yahoo.co.in

Disclaimer: IJAPR is solely owned by Mahadev Publications - dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IJAPR cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of IJAPR editor or editorial board members.