



Research Article

EFFICACY OF PICHHA BASTI IN ULCERATIVE COLITIS: AN OBSERVATIONAL STUDY

Chiterlekha^{1*}, Swati S. Deshpande², Baidyanath Mishra³

*1PG Scholar, ²Guide and Professor and Head, Dept. of PG Studies in Panchakarma, SKAMCH & RC, Bangalore, Karnataka, India.

³Professor and Head, Department of Rachana Sharira, Dy. Dean-PG Studies, Shri Kalabyraveshwaraswamy Ayurvedic Medical College Hospital & Research Centre Vijayanagar, Bangalore, India.

Received on: 27/11/2014

Revised on: 12/12/2014

Accepted on: 25/12/2014

ABSTRACT

Ulcerative colitis is a chronic idiopathic inflammatory bowel disease with a relapsing nature. It is a very challenging disease affecting a patient during the most active period of his life i.e. 20 to 40 years of age. The main features are ano-rectal bleeding with increased frequency of bowel evacuation, general debility and abnormal structural pathology in the descending colon, particularly sigmoid colon. In modern medical science, there is no permanent curative treatment for this disease. This study can be helpful for reducing the need of steroids and surgical processes in the patients of ulcerative colitis. A clinical study of 5 patients of ulcerative colitis has been conducted at the O.P.D. and I.P.D. SKAMC & RH, Bangalore. They were treated with *Pichha basti* for 8 days in *Yoga basti* schedule. Results were analyzed statistically by using the 't' test. In this study, it was observed that the symptoms and signs were reduced by more than 73% with a highly significant result.

KEYWORDS: Ulcerative colitis, *Pichha basti*.

INTRODUCTION

Ulcerative colitis (UC) and Crohn's disease (CD) are chronic inflammatory bowel diseases which pursue a protracted relapsing and remitting course, usually extending over years.¹ Ulcerative colitis, a disorder in which inflammation affects the mucosa and submucosa of the colon.

The rapid changes in incidence and prevalence in Asia may point to environmental changes associated with the 'Westernisation' of lifestyle (e.g. dietary changes, smoking, etc.) as potential risk factors². Its incidence is more during 20 to 40 years of age and men are slightly more likely to be affected with UC.

The major symptoms of UC are diarrhoea, rectal bleeding, tenesmus, passage of mucus, and crampy abdominal pain³. The severity of symptoms correlates with the extent of disease. Although UC can present acutely, symptoms usually have been present for weeks to months. Occasionally, diarrhoea and bleeding are so intermittent and mild that the patient

does not seek medical attention. The first attack is usually the most severe and thereafter the disease is followed by relapses and remissions. Ulcerative colitis can be associated with complications like massive haemorrhage, toxic mega colon, perforation, stricture and colo-rectal carcinoma.

Ulcerative colitis only involves the colon, particularly the sigmoid colon. According to *Ayurveda*, *Pakvashya* and *Sthula guda* are the *mulasthanas* for *Purishavaha srotas*⁴. So this disease can be considered under *Purishavaha srotas* and can be correlated with *Raktatisra*. *Pichha basti* is mentioned as a management of *Raktatisara* in *Charaka Samhita*⁵.

AIMS AND OBJECTIVES

To evaluate the efficacy of *Pichha basti* in the management of ulcerative colitis.

MATERIALS & METHODS

The present study is an observational study with open label study design. All subjects

were administered *Pichha basti* for a period of 8 days. *Pichha basti* is prepared as per the description in *Charaka samhita chikitsa sthana*. Formulated products purchased from the approved pharmacy stores manufactured by approved pharmaceutical manufacturers.

Selection of patient: An open, randomized, clinical study was conducted on 5 patients diagnosed as Ulcerative colitis and selected from OPD & IPD of SKAMC & RH, Bangalore.

Subjective Parameter

- Bowel frequency
- Blood in stool
- Abdominal pain
- Weakness

Objective Parameter

- Red blood cells in stool
- Pus cells in stool

Assessment criteria

Improvements in the patients were assessed on the basis of relief in the subjective and objective parameters. All the subjective and objective parameters were scored and assessed before as well as after the treatment.

Gradation of subjective & objective parameters

1. Bowel frequency

- 0 = 1 or 2 times in a day
- 1 = 3 or 4 times in a day
- 2 = 5 or 7 times in a day
- 3 = 8 or 12 times in a day
- 4 = More than 12 times in a day

2. Blood in stool

- 0 = No bleeding
- 1 = Occasional bleeding in stool (not daily)
- 2 = Bleeding daily but less than 4 times per day
- 3 = Bleeding daily but less than 8 times per day
- 4 = Bleeding daily more than 8 times per day

3. Abdominal pain

- 0 = No pain
- 1 = Tolerable pain
- 2 = Mild pain
- 3 = Moderate pain
- 4 = Severe pain

4. Weakness

- 0 = No weakness
- 1 = Tolerable weakness
- 2 = Mild weakness
- 3 = Moderate weakness
- 4 = Severe weakness

5. Red blood cells in stool

- 0 = No RBC in stool
- 1 = Less than 15/hpf
- 2 = 16-30/hpf
- 3 = 31-50/hpf
- 4 = Plenty or more than 50/hpf

6. Pus cells in stool

- 0 = No pus cells in stool
- 1 = Pus cell Less than 15/hpf
- 2 = Pus cell 16-30/hpf
- 3 = Pus cell 31-50/hpf
- 4 = Plenty or more than 50/hpf Pus cell

Sample Size and Method

Sample size: 5 without any drop outs.

Study design: Open level clinical trial

Number of groups: Single arm

Drug schedule and Dosage

<i>Niruha basti</i>	<i>Anuvasna basti</i>
<i>Madhu</i> = 100ml	<i>Jatayadi taila</i> 60ml with a pinch of <i>Saindhava</i>
<i>Saindhava</i> = 6gms	
<i>Sneha</i> = <i>Jatayadi taila</i> = 60ml + <i>Changeari gritha</i> = 60ml	
<i>Kalka</i> = <i>Yastimadhu choorna</i> (20gm) + <i>Pushyanuga choorna</i> (20gm)	
<i>Kwatha</i> = <i>Shalmalii twak</i> + <i>Yastimadhu sidhha ksheera paka</i> = 240ml	

Procedure

Poorva karma - *Sthanika abhayanga* with *Moorchita tila taila* and *Nadi sweda*.

Pradhana karma - Patient was made to lie in left lateral position for administration of *Basti*. *Niruha basti* was given with empty stomach and *Anuvasana basti* after the food.

Paschata karma - lift the legs, patting to the buttocks, and anti-clockwise massage to abdomen.

Duration of Treatment: = 8 days in *Yoga basti* schedule.

Inclusion criteria

- Diagnosed cases of ulcerative colitis
- Patient who are fit for *Basti karma*

Exclusion criteria

- Patients who had undergone surgeries

- Patients who had complications like anal fistula etc.

Investigations: Stool examination

OBSERVATIONS & RESULT


Out of 5 patients, 1 (20%) patient was female and 4 (80%) patients were male; all the patient (100%) were between the age group of 25-30, with regards to addictions, 3 (60%) patients were non smoker, 2 (40%) patients were ex-smoker and all the 5 (100%) patients

were used to take spicy food. 3 (60%) patients belonged to the 1 to 2 years of chronicity, 2 (40%) patients were found in 3 to 5 years of chronicity group.

Bowel frequency was reduced by 81%, Bleeding in stool was reduced by 80%, Abdominal pain was reduced by 87%, Weakness was reduced by 73%, RBC in stool was decreased by 85%, Pus cells in stool were decreased by 83%. These all parameters showed statistically highly significant result.

Table 2: Changes in before treatment & after treatment (n=5)

Parameter	Mean		Improvement (%)	SD	SE	t-value	P-value
	BT	AT					
Bowel frequency	3.2	0.8	81%	0.5477	0.2739	10.61	<0.001
Blood in stool	3	0.4	80%	0.5477	0.2739	9.798	<0.001
Abdominal pain	3	0.4	87%	0.5477	0.2739	9.494	<0.001
Weakness	2.8	0.4	73%	0.4472	0.2	9	<0.001
Red blood cells in stool	2.4	0.4	85%	0.4472	0.2	11	<0.001
Pus cells in stool	2.4	0.4	83%	0.7071	0.3162	6.325	<0.001



Graph: Shows improvement in the treatment in before treatment and after treatment

DISCUSSION

Ulcerative colitis is a mucosal disease that usually involves the rectum and extends proximally to involve all part of colon. The cause is unknown, but several theories have been put forward of which the main ones are infective, nutritional, psychosomatic & immunological. According to *Ayurveda*, we can consider this disease under *Purisha vaha srotas* due to involvement of mainly *Sthula guda* and its treatment was explained as *Atisara chikitsa*.

Basti is the best mode of administration of the drug as *Acharya Charaka* has quoted 'the medicine administered through anus is more effective in the disorder of *Pakvashya*. In *Vata* or *Vatapitta pradhana atisara*, *Basti* is indicated to overcome the disease and at the same time it gives *Bala* to *Kati*, *Pakvashya sthana*⁶.

According to *Sushruta samhita*, *Taila avcharana* in the form of *Pichu*, *Seka* & *Anuvasana basti* is indicated in *Guda dourbalaya*. *Pichha basti* is indicated in *Alpa-alpa*, or *Bahu raktapravruti*, *Sashoola*, *Vayu vibdhata* like conditions⁷. Especially when *Ruja lakshana* is dominant then mainly *Pichha basti* is indicated⁸.

In this present study, the drugs used for *Pichha basti* are *Madhu*, *Saindhava*, *Jatayadi Taila*, *Changeri Ghrita*, *Yashtimadhu Choorna*, *Pushyanuga Choorna* for *Kalka* and *Shalmali twak* with *Yashtimadhu siddha ksheera paka* for *Kwatha* and *Jatyadi taila* for *Anuvasana basti*.

Madhu is *Sheeta veerya* and *Madhura rasa*, *Kashaya anurasa* with *Shodhana*, *Ropana* properties. So its *Ropana* property helps to heal the ulcer. *Jatyadi taila (JT)*, by its best *Vranaropaka* property heals the ulceration and

its *Raktastambhana* property provides haemostasis. The phytochemical evaluation of *Jatyadi taila* has revealed presence of flavonoids, essential oils, tannins, glycosides, steroids and alkaloids. Tannins & phytosterols promote the wound healing process with increased capillary formation & fibroblast proliferation enhancing the rate of epithelisation.

The drugs in *Changeri ghrta* are having *Shothahara*, *Vranaropaka* and *Raktasthabhaka* properties which are helpful in this disease. *Yastimadhu* is having properties like *Vatapittajit*, *Raktaprasadana*, *Balya*, *Varnya* and *Madhura rasa*, *Madhura vipaka* with *Sheeta veerya*. Due to presence of Glycyrrhizin, this is a gastroprotective and increases the rate of mucus secretion by the gastric mucosa. The anti-allergic properties of *Yashtimadhu* have been attributed to the corticosteroid-like activity of glycyrrhizin and glycyrrhetic acid, also known as enoxolone. *Pushyanug Churna* is potent as antibacterial and is mainly indicated in *Rakta atisara* and *Rakta arshas*.

For *Kwatha* purpose, *Shalmali* is used which is having *Madhura rasa*, *Madhura vipaka*, *Sheeta veerya*, *Snigdha*, *Pichhala guna*. *Yashtimadhu siddha ksheera paka* in that *Ksheera* which contains *Madhura rasa*, *Sheeta veerya*, *Pichhil guna* and indicated in *Vata*, *Pitta*, *Rakta atisara*. So using *Ksheera paka* with *Yashtimadhu* is more effective.

So the drugs used in *Basti Karma* are having combined effect like *Vata pitta shamana*, *Rakta sthambhana*, *Vrana shodhana* and *Vrana ropana*.

CONCLUSION

Basti karma has shown improvement or cure of the ulcerations of the bowel by enhanced healing and subsiding inflammation & irritability of the colon. It also improves functions of *Apana Vayu*, situated in *Pakwashaya* and results into improvement of ulcerations and regulation of evacuation of the

colon. In the present study the mean scores of frequency of diarrhoea, blood and mucous in stool, pain abdomen, weakness, pus cell in stool are compared to before and after treatment is highly significant. Keeping in mind the complications of surgical management, this *Pichha basti* is definitely a better option. Although the number of patients included in this study was small, but this study provides a basis for further evaluation by a large longitudinal study to substantiate its findings.

ACKNOWLEDGEMENTS

Dr. N.Anjaneya Murthy, Dean, Dept of P.G studies, SKAMCH & RC.

Dr. Kiran.M.Goud, Principal, SKAMCH & RC, Vijayanagar, Bangalore.

REFERENC ES

1. Sir Stanley Davidson, Davidson's Principal and practice of medicine, edited by Nicholas A.Boon, 20th edition, reprint 2008, pp - 1381, pg - 910.
2. API text book of medicine, edited by YP Munjal, 9th edition, pp- 2165, pg-829.
3. Fauci, Braunwald, Kasper, Hauser, Longo, Jameson - Harrison's Principles of Internal medicine, New York, McGraw-Hill Medical Publishing Division, 18th Edition - 2008, Volume -II, pp - 2754, pg - 2481.
4. Agnivesha, Charaka Samhita, Ayurveda Deepika Commentary of Chakrapani, edited by; Vaidya Yadavji Trikrumji Acharya, Choukambha Surabharati Prakashan, Varanasi, reprint-2011, pp - 738, pg - 251.
5. Ibid, pp - 738, pg - 552.
6. Ibid, pp - 738, pg - 553.
7. Sushruta, Sushruta Samhita, Nibandha Samgraha Commentary of Sri Dalhanacharya and Nyaya Chandrika Panjika on Nidanasthana Commentary of Sri Gayadasacharya, by; Vaidya Yadavji Trikrumji Acharya, Choukambha Surabharati Prakashan, Varanasi, reprint-2008, pp - 824, pg - 704.
8. Ibid, pp - 824, pg - 705.

Cite this article as:

Chiterlekha, Swati S. Deshpande, Baidyanath Mishra. Efficacy of Pichha Basti in Ulcerative Colitis: An Observational Study. Int. J. Ayur. Pharma Research. 2014;2(8):23-26.

Source of support: Nil, Conflict of interest: None Declared

*Address for correspondence

Dr. Chiterlekha

3rd year PG Scholar

Dept, of Panchakarma, SKAMCH & RC Vijayanagar Bangalore

Karnataka, India.

Email: dr.chiterlekha@gmail.com