Int. J. Ayur. Pharma Research, 2014; 2(4): 103-106 ISSN: 2322 - 0910



International Journal of Ayurveda and Pharma Research

## **Research Article**

# INTEGRATED APPROACH IN THE MANAGEMENT OF BRONCHIAL ASTHMA: A RETROSPECTIVE ANALYSIS

## Memane N1\*, Shanke A1

<sup>1</sup> Departments of Ayurveda and Pediatrics, Bhakti Vedanta Hospital and Research Institute, Srishti Complex, Mira Road East, Thane, Maharashtra, India.

Received on: 14/07/2014	Revised on: 07/08/2014	Accepted on: 26/08/2014
-------------------------	------------------------	-------------------------

#### ABSTRACT

Bronchial asthma is a widely occurring respiratory tract disorder with a poor quality of life, associated with adverse reactions and many achieve incomplete cure. Recently, an integrated approach of allopathy with lifestyle management, psychosocial approaches and evidence based complementary medicine has been promoted in the management of bronchial asthma. Hence, the present study was undertaken to evaluate the same.

**Methods:** The present study was a retrospective analysis of patients diagnosed as asthmatics who were being managed through integrated approach that includes a combination of ayurvedic drugs, lifestyle management and the allopathic agents. The ayurvedic drugs include the respiratory tonics and naturally occurring bronchodilators such as *Shwaskaschintamani, Shwaskuthar ras, Vasakasavam, Vasavleham, Mahalaxmivilas rasa.* The allopathic drugs include short and long-acting beta-2 agonists and inhalational corticosteroids depending on the intensity of bronchial asthma.

**Results:** A total of 36 children and 71 adults were identified as asthmatics with a median (range) age (in years) of 8 (2-13) and 46 (18-64). A significant resolution of all the symptoms except for hyperacidity was observed at all the follow-up periods as compared to the baseline (P < 0.05).

**Conclusion:** A holistic approach of adding alternative therapies to the conventional pharmacotherapy offers the asthmatic patients a potential to safely control their symptoms in Indian patients.

**KEYWORDS:** Ayurveda, Bronchial asthma, Polyherbal preparations, Complementary and Alternative medicine.

#### **INTRODUCTION**

Bronchial asthma is a widely occurring respiratory tract disorder with a prevalence of 5-15% and a total of 15-20 million asthmatics in the country.<sup>[1,2]</sup> A poor quality of life has been reported in individuals with bronchial asthma in India.<sup>[3]</sup> The parents of asthmatic children have also been found to have a poor quality of life.<sup>[4]</sup> Despite the presence of many effective allopathic medicines for managing the disease, their side effects are of major concern especially when used for long-term.<sup>[5]</sup> Hence, recently an integrated approach of allopathy with lifestyle management, psychosocial approaches and evidence based complementary medicine has been promoted in the management of bronchial asthma.<sup>[6]</sup> Our hospital is a tertiary care centre and we recently started implementing this integrated approach for treating bronchial asthma. Considering its infancy stage, hardly any studies have been performed to determine the usefulness of integrated approach, especially in Indian set up. Hence, we did a retrospective analysis of drug-naive patients with bronchial asthma who underwent management with the integrated approach in our institute.

#### METHODS

The present study was a retrospective analysis of those patients diagnosed to have bronchial asthma who were being managed through integrated approach between December 2013 and June 2014 in the Ayurveda outpatient department in Bhakti Vedanta Hospital and Research Institute. The study was commenced after obtaining institutional ethics committee approval and a waiver was obtained for informed consent. The integrated approach of management includes the combination of ayurvedic drugs, lifestyle management and allopathic agents. The ayurvedic drugs include the respiratory tonics and naturally occurring bronchodilators such as Shwaskaschintamani. Shwaskuthar ras, Mahalaxmivilas rasa (1 gram of Rasa medicines for 1 month in divided doses), Vasakasavam (1-10 ml twice a day) and Vasavleham (2-10 gram twice a day). As per subjective treatment, *Prakriti abd dosha* imbalance, any one or all medicines were used. Treatment modification (dose changes) was done on age but also was based on Dushya (Tissue affected), Desham, Bala (Immunity), Kalam (Time or seasonal change), Analam (Agni digestive fire), *Prakriti* (Nature). The allopathic drugs include short and long-acting beta-2 agonists and inhalational corticosteroids depending on the intensity of bronchial asthma. The lifestyle changes include standard dietary patterns to be followed (avoid fermented, stored, cold food items) and a regular sleep pattern. The presenting symptoms were classified into various categories such as wheeze, cough,

dyspnoea, sleep disturbance, hyperacidity and cold. All the patients who underwent integrated therapy were asked to follow up at 15 days, 1 month and 3 months following the initiation of therapy. Patients were classified into children (aged < 12 years) and adults (12 years and above). Descriptive statistics was used for analyzing the demographic details and Cochran Q test was used for analyzing the significance of resolution of each of the presenting symptoms in these patient categories. Normality of the continuous variable was ascertained using Kolmogorov-Smirnov test and non-normal data were represented using median (range).

## RESULTS

A total of 36 children and 71 adults were identified as asthmatics with a median (range) age (in years) of 8 (2-13) and 46 (18-64). Table 1 and 2 depict the total number of individuals with various symptoms of asthma and their resolution at the follow-up periods in paediatric and adult population. A significant resolution of all the symptoms except for hyperacidity was observed at all the follow-up periods as compared to the baseline (P < 0.05).

8		
Table 1: Number of child	re <mark>n</mark> with each symptom	at different time points

Symptoms	Baseline	First follow-up (15 days)	Second follow- up (1 month)	Third follow-up (3 months)
Wheeze*	36	1	1	1
Cough*	13	7 UAPR	1	1
Dyspnea*	9	3	1	0
Disturbance of sleep*	4	1	1	0
Hyperacidity	2	1	1	0
Cold*	10	5	1	1

\* - P < 0.05 (significant)

Symptoms	Baseline	First follow-up (15 days)	Second follow- up (1 month)	Third follow-up (3 months)
Wheeze*	29	0	0	0
Cough*	79	14	2	0
Dyspnea*	33	8	3	0
Disturbance of sleep*	49	2	0	0
Hyperacidity	2	0	0	0
Cold*	71	8	2	0

\* - P < 0.05 (significant)

## DISCUSSION

The present study was a retrospective analysis of patients diagnosed to have bronchial asthma who were managed through integrated approach in a tertiary care hospital. We found a significant improvement of all the patients with this therapy.

Bronchial asthma is a widely prevalent respiratory tract disorder with limited options available in the allopathic system of medicine. The WHO has estimated that 16 million disability-adjusted life-years are lost annually due to asthma, representing 1% of the total global disease burden. It is a public health problem not just for high-income countries; it occurs in all countries regardless of the level of development.<sup>[7]</sup> A recent study by Kotwani *et al.* revealed the presence of wide gaps between the desirable and the actual management-practices on asthma in Indian set up.<sup>[8]</sup> The authors found an unsatisfactory patient-doctor contact, as well as poor knowledge about asthma and its management amongst patients. A cross-sectional study from an Indian metropolitan city revealed that majority of the asthmatic patients were administered short-acting beta-2 agonists (100%) and anticholinergics (80%) followed by corticosteroids inhalational (71%) and theophylline (5%).<sup>[9]</sup> Around 15% of the individuals experienced adverse effects in the form of palpitation, tremors, vomiting and dry mouth to the allopathic medicines. Hence, there is a need for using alternate systems medicinal systems for managing bronchial asthma.

Ayurveda, the great indigenous system of medicine of India refers to bronchial asthma as Tamaka Swasa and contributes several modalities of treatment for the same.<sup>[10]</sup> Although individual herbs have shown good amount of efficacy, polyherbal combinations are found to be well-accepted, safe and effective in asthma.<sup>[11]</sup> Polyherbal preparations cause target the multiple components of the disease pathway. Manv avurvedic such agents as Shwaskaschintamani. Shwaskuthar ras Vasakasavam, Vasavleham, Mahalaxmivilas rasa, Nidigdhikadi kvatha and Padmapatradi yoga have been shown to be effective in the control of asthmatic attacks.<sup>[12,13]</sup> In the same study, it was also observed that around 17% of the individuals were also using alternative medicine. In the present study we have carried out an integrated approach of combining both the ayurvedic agents with allopathic system of medicine along with lifestyle alterations. Despite inadequate literature on the possible interactions between the ayurvedic drugs used in the study with allopathic agents, we did not observe any adverse drug reactions. Although there were no previous studies with such approach of allopathic drugs with ayurvedic agents, a study combining yoga with ayurvedic drugs for treating bronchial asthma has shown a significant improvement in the ventilation parameters.<sup>[14]</sup>

This is the first study reporting the usefulness of an integrated approach for managing bronchial asthma from India to the best of our knowledge. However, the study is also limited in being a retrospective analysis, small number of study participants, and absence of any control group and objective measures of the disease without quality of life of the patients. The findings of this study have to be confirmed in large population based studies.

To conclude, a holistic approach adding alternative therapies to the conventional pharmacotherapy offers the asthmatic patients a potential to safely control their symptoms in Indian patients.

## ACKNOWLEDGEMENT

We thank Medical Research Department of Bhakti Vedanta Hospital and Research Institute for rendering help in the preparation of manuscript.

## REFERENCES

1. Bronchial asthma. WHO fact sheet. Available at:

http://www.who.int/mediacentre/factshe ets/fs206/en (Accessed on 08 July 2014).

- 2. Pal R, Dalal S, Pal S. Prevalence of bronchial asthma in Indian children. Ind J Comm. Med 2009;34:310-6.
- 3. Singh M, Mathew JL, Malhi P, Naidu AN, Kumar L. Evaluation of quality of life in Indian children with bronchial asthma using a disease-specific and locally appropriate questionnaire. Acta Paediatr. 2004;93:554–5.
- 4. Kumari V, Gupta V, Piplani S, Bhatia BD, Upadhayay SK. Parental stress and coping techniques in parents of children with bronchial asthma. Indian J Allergy Asthma Immunol 2011;25:103-8.
- Sadhna K, Abbulu K. Holistic approach to management of asthma. IJRAP 2010;1:367-83.
- 6. Hassed C. An integrative approach to asthma. Australian Family Physician 2005;34:573-6.
- World Health Organization. Asthma fact sheet. Available at: <u>http://www.who.int/mediacentre/factshe</u> <u>ets/fs307/en/</u> (Accessed on 09 July 2014).
- 8. Kotwani A, Chhabra SK, Tayal V, Vijayan VK. Quality of asthma management in an

urban community in Delhi, India. Indian J Med Res 2012;135:184-92.

- 9. Kamath L, Kulkarni C. Study of pattern of drug treatment in patients with exacerbations of bronchial asthma in an emergency ward of a teaching hospital: A prospective, cross-sectional study. Journal of Chemical and Pharmaceutical Research 2012;4:1815-21.
- 10. Sharma PV. 1st ed. Varanasi: Chaukhamba Orientalia; 1982. Charaka Samhita-Shvasa Chikitsa- English translation. Chapter 17 Sloka 45-76.
- 11. Ng TP, Wong ML, Hong CY, Koh KT, Goh LG. The use of complementary and alternative

## Cite this article as:

Memane N, Shanke A. Integrated Approach in the Management of Bronchial Asthma: A Retrospective Analysis. Int. J. Ayur. Pharma Research. 2014;2(4):103-106. *Source of support: Nil, Conflict of interest: None Declared* 



medicine by asthma patients. QJM. 2003; 96:747–54.

- 12. Savita C, Reddy GR, Jayant H, Prerna J. A clinical study on efficacy of Nidigdhikadi kvatha on Tamaka Shvasa (Bronchial asthma). International Research Journal of Pharmacy. 2013;4:141-4.
- Panda AK, Doddanagali SR. Clinical efficacy of herbal Padmapatradi yoga in bronchial asthma. J Ayurveda Integr Med. 2011;2:85-90.
- 14. Rao YC, Kadam A, Jagannathan A, Babina N, Rao R, Nagendra HR. Efficacy of naturopathy and yoga in bronchial asthma. Indian J Physiol Pharmacol 2014;58:232-38.

\*Address for correspondence Dr. Nanasaheb Memane Consultant Ayurvedic physician Bhakti Vedanta Hospital and Research Institute, Srishti Complex, BhaktiVedanta Swami Marg Mira Road East, Thane Maharashtra, India. Email: apsankhe@bhaktivedantahospital.com