ISSN: 2322 - 0902 (P)
ISSN: 2322 - 0910 (O)

International Journal of Ayurveda and Pharma Research

Research Article

INTEGRATIVE MIND-BODY THERAPIES FOR NON-ALCOHOLIC FATTY LIVER DISEASE: A PRE-POST INTERVENTIONAL STUDY AMONG SECONDARY SCHOOL TEACHERS

S. Venkatesan[1]*, D R Rajendran[2]

*[1]Research Scholar, Manipur International University, Manipur

[2]Professor, Sandip Institute of Technology and Research, Nasik, Maharashtra, India.

Article info

Article History:

Received: 09-06-2026

Accepted: 06-07-2026

Published: 15-08-2026

KEYWORDS:

Non-Alcoholic Fatty Liver Disease, Mind-Body Therapies, Yoga, Raja Yoga Meditation, Varma Therapy, Occupational Health, Lifestyle Medicine, Teachers.

ABSTRACT

Non-Alcoholic Fatty Liver Disease (NAFLD) affects approximately 25% of the global adult population, with teachers representing a particularly vulnerable group due to sedentary work patterns, irregular routines, and chronic occupational stress. Integrative mind-body therapies may offer multifaceted benefits addressing both physiological and psychological determinants of NAFLD. This study evaluated the effectiveness of an integrated intervention combining Raja Yoga Meditation, Yoga, Varma therapy, dietary counselling, and lifestyle modification among secondary school teachers diagnosed with Grade I or Grade II NAFLD. A pre-test and post-test experimental design was employed with 30 participants (25-60 years) recruited through purposive sampling. Baseline and post-intervention assessments included liver function tests (ALT, AST), ultrasonography grading, body mass index, waist circumference, Perceived Stress Scale-10, and WHOQOL-BREF. The 8-week structured intervention comprised daily Raja Yoga meditation (30 min), daily Yoga practice (45 min), Varma therapy (three times/week), weekly dietary counselling, and lifestyle modification recommendations. Data were analyzed using paired t-tests, with significance set at $p<0.05$. Post-intervention analysis revealed statistically significant reductions in ALT (54.6±22.3 to 34.2±14.8 U/L, $p<0.001$), AST (46.8±18.5 to 30.6±12.1 U/L, $p<0.001$), BMI (28.4±4.2 to 26.8±3.9kg/m^2, $p<0.001$), and waist circumference (94.3±10.8 to 90.1±10.2 cm, $p<0.001$). Perceived stress scores decreased significantly (24.6±6.8 to 16.4±5.2, $p<0.001$), while quality of life improved across all domains ($p<0.001$). Ultrasonographic assessment showed significant improvement, with 83.3% of participants demonstrating regression in fatty liver grade. The integrated mind-body intervention produced significant improvements in hepatic, anthropometric, and psychosocial outcomes among teachers with NAFLD, suggesting that such approaches may serve as effective complementary strategies for managing NAFLD in high-stress occupational populations. Future randomized controlled trials with larger samples and longer follow-up are warranted.

INTRODUCTION

Non-Alcoholic Fatty Liver Disease (NAFLD) constitutes a spectrum of liver disorders characterized by excessive triglyceride accumulation within hepatocytes, occurring in the absence of significant alcohol consumption. The pathological continuum ranges from simple hepatic steatosis to non-alcoholic steatohepatitis (NASH), fibrosis, cirrhosis, and hepatocellular carcinoma. NAFLD is strongly associated with metabolic syndrome components and is widely recognized as the hepatic manifestation of insulin resistance.

The global prevalence of NAFLD has increased dramatically over the past two decades, driven by dietary changes, declining physical activity, urbanization, and escalating psychological stress. Current epidemiological estimates indicate that approximately 25% of the global adult population is affected, positioning it as one of the leading causes of chronic liver disease worldwide.

Access this article online

Quick Response Code

https://doi.org/10.47070/ijapr.v14i8.4287

Published by Mahadev Publications (Regd.)
publication licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0)

Teachers constitute an occupational group facing unique physical and psychological challenges that predispose them to metabolic disorders. Extended teaching hours, administrative responsibilities, classroom management demands, academic pressures, and limited opportunities for self-care frequently contribute to chronic stress and maladaptive lifestyle practices. These factors promote obesity, dyslipidemia, insulin resistance, and ultimately, fatty liver disease.

The cornerstone of NAFLD management remains lifestyle modification, encompassing weight reduction, dietary regulation, and increased physical activity. However, adherence to conventional recommendations is frequently suboptimal due to psychological barriers, motivational challenges, and competing occupational demands. This has generated growing interest in holistic interventions that simultaneously address physical, mental, and emotional dimensions of health.

Mind-body therapies, including meditation, Yoga, and traditional therapeutic modalities, offer promising adjunctive approaches. Studies on Yoga have demonstrated improvements in body mass index, waist circumference, blood glucose, lipid profiles, and stress-related outcomes. Meditation practices have been associated with enhanced emotional regulation, reduced anxiety, lowered cortisol secretion, and improved autonomic balance. "Raja Yoga Meditation, specifically, has shown beneficial effects on autonomic functions and psychological well-being (Sharma et al., 2015)."

Despite the increasing prevalence of NAFLD among working professionals and the theoretical benefits of integrative approaches, scientific studies evaluating the combined effectiveness of Raja Yoga Meditation, Yoga, and Varma therapy in reversing NAFLD among teachers remain limited. This study addresses this gap by systematically evaluating an integrated intervention in a vulnerable occupational population.

Literature Review

Global Burden and Epidemiology of NAFLD

Younossi et al. (2016) conducted a comprehensive meta-analytic assessment establishing NAFLD as the most prevalent chronic liver disease worldwide, with strong associations with obesity and metabolic syndrome. Their analysis revealed that lifestyle modification remains the first-line treatment and significantly improves hepatic steatosis. Subsequent research has highlighted the escalating global burden and the imperative for effective non-pharmacological interventions.

Lifestyle Interventions in NAFLD Management

Kenneally, Sier, and Moore (2017) systematically reviewed dietary and physical activity interventions in NAFLD, concluding that lifestyle modification constitutes the cornerstone of treatment. Their findings demonstrated that adopting a healthy diet, engaging in regular physical activity, and achieving gradual weight loss significantly reduces liver fat accumulation. Even modest weight reduction (3-5%) improves hepatic steatosis, insulin sensitivity, and inflammatory markers associated with NAFLD.

Paradigm Shift: MAFLD Framework

Eslam et al. (2020) introduced the concept of Metabolic Dysfunction-Associated Fatty Liver Disease (MAFLD) to more accurately reflect underlying etiological factors. They proposed that metabolic dysfunction, rather than merely the absence of alcohol consumption, should constitute the primary diagnostic criterion. The MAFLD definition encompasses patients with obesity, type 2 diabetes, or other metabolic abnormalities, enhancing clinical relevance and facilitating improved patient identification and management.

Yoga and Mind-Body Interventions

Pascoe and Bauer (2015) systematically reviewed randomized controlled trials on yoga's effects on stress measures and mood, demonstrating significant improvements across multiple psychological outcomes. Cramer et al. (2016) found that yoga improves health-related quality of life and mental health outcomes. Regarding NAFLD specifically, yoga interventions have shown improvements in body mass index, waist circumference, blood glucose, lipid profiles, and stress-related outcomes.

Goyal et al. (2014) conducted a systematic review and meta-analysis of meditation programs, establishing significant effects on psychological stress and well-being. Black and Slavich (2016) provided evidence that mindfulness meditation influences immune system function through stress reduction pathways. "Raja Yoga Meditation specifically has demonstrated beneficial effects on autonomic functions, including improved heart rate variability and reduced sympathetic activation (Sharma et al., 2015)."

Teacher Stress and Health Outcomes

Kyriacou (2001) provided foundational research on teacher stress, identifying key occupational stressors and their health implications. Jennings and Greenberg (2009) established connections between teacher social-emotional competence, classroom outcomes, and occupational health. Roeser et al. (2013) demonstrated that mindfulness training improves teacher well-being and professional development. These studies establish teachers as an occupational group at elevated risk for stress-related health conditions.

OBJECTIVES AND HYPOTHESES

Primary Objective

To evaluate the effectiveness of an integrated intervention comprising Raja Yoga Meditation, Yoga, Varma therapy, dietary counselling, and lifestyle modification in reversing Non-Alcoholic Fatty Liver Disease among secondary school teachers.

Secondary Objectives

1. To assess changes in liver function parameters (ALT, AST).
2. To evaluate reduction in ultrasonographic fatty liver grading.
3. To assess changes in anthropometric parameters (BMI, waist circumference).
4. To evaluate stress reduction using Perceived Stress Scale-10.
5. To assess improvements in quality of life using WHOQOL-BREF.

Hypotheses

Null Hypothesis (H_0): The integrated intervention will not produce significant effects on liver function parameters, anthropometric measurements, stress levels, quality of life, or fatty liver grade among teachers with NAFLD.

Alternative Hypothesis (H_1): The integrated intervention will produce significant improvements in liver function parameters, anthropometric measurements, stress levels, quality of life, and fatty liver grade among teachers with NAFLD.

METHODOLOGY

Study Design

The study employed a pre-test and post-test quasi-experimental design (one-group pretest-posttest design). Participants served as their own controls, with baseline assessments compared to post-intervention outcomes following an 8-week integrated intervention. This design was selected for feasibility given the exploratory nature of the study and the practical constraints of recruiting teachers with NAFLD for a controlled trial.

Study Setting and Duration

The study was conducted between January and April 2026 at a wellness centre in Chennai city, India. Educational institutions and naturopathy wellness centres served as recruitment sites. The total study duration was 12 weeks, comprising recruitment and baseline assessment (2 weeks), intervention period (8 weeks), and post-intervention assessment (2 weeks).

Participant Selection

Sampling Technique

Purposive sampling was employed to recruit 30 secondary school teachers diagnosed with Grade I or Grade II NAFLD. This approach was appropriate given the specific population and condition under investigation.

Inclusion Criteria

- Secondary school teachers aged 25 to 60 years.
- Diagnosed with Grade I or Grade II NAFLD confirmed by ultrasonography.
- Willing to participate and provide written informed consent.
- Available to attend the 8-week intervention program.
- No prior regular practice of Yoga or meditation.

Exclusion Criteria

- Alcoholic liver disease (self-reported or confirmed by medical history).
- Viral hepatitis (Hepatitis B or C positive).
- Cirrhosis or hepatocellular carcinoma.
- Severe systemic illnesses (cardiac, renal, or respiratory failure).
- Pregnancy or lactation
- Regular practice of Yoga or meditation in the preceding 6 months.
- Participation in other clinical trials.

Assessments and Outcome Measures

Primary Outcome Measures

Liver Function Tests: Serum ALT and AST assessed using standardized biochemical assays (fully automated clinical chemistry analyser) at baseline and post-intervention.

Ultrasonography Assessment: Fatty liver grading (Grade I, II, III) performed by experienced radiologists using standardized protocols. Grading criteria: Grade I (mild steatosis), Grade II (moderate steatosis), Grade III (severe steatosis).

Secondary Outcome Measures

Anthropometric Measurements: BMI calculated as weight (kg)/height $(m)^2$. Waist circumference measured at the midpoint between the iliac crest and lower costal margin using standardized technique.

Perceived Stress: Perceived Stress Scale-10 (PSS-10), a 10-item validated questionnaire assessing global perceived stress. Score range: 0-40; higher scores indicate greater perceived stress.

Quality of Life: WHOQOL-BREF, a 26-item questionnaire assessing physical, psychological, social, and environmental domains. Higher domain scores indicate better quality of life.

Intervention Protocol

Raja Yoga Meditation Component

Participants practiced Raja Yoga meditation daily for 30 minutes under trained guidance. Sessions included:

- Positive thought generation (10 min)
- Self-awareness exercises (5 min)
- Visualization techniques (5 min)

- Soul-consciousness practices (5 min)
- Silent reflection (5 min)

Participants received a structured meditation manual and were encouraged to practice independently in addition to guided sessions.

Yoga Component

Yoga practice comprised 45 minutes daily, incorporating:

Preparatory/Loosening Exercises (10 min): Neck rotations, shoulder rotations, arm swings, trunk rotations, knee movements, ankle rotations.

Asanas (20 min): Surya Namaskar (6 rounds), Bhujangasana, Dhanurasana, Ardha Matsyendrasana, Pavanamuktasana, Vajrasana, Paschimottanasana, Trikonasana, Setu Bandhasana.

Pranayama (10 min): Kapalabhati (60 strokes, 3 rounds), Anuloma Viloma (10 rounds), Deep Diaphragmatic Breathing (5 min), Bhramari (5 rounds).

Relaxation (5 min): Shavasana, Yoga Nidra.

Varma Therapy Component

Varma therapy was administered three times per week by trained practitioners. Selected Varma points included:

- Periyaathi suruki (thoracic region, circular massage, 2 min)
- Palla varmam (upper lumbar, deep pressure, 2 min)
- Siriyaathi suruki (mid-thoracic, gentle stimulation, 2 min)
- Adappa Kalam (abdominal, clockwise massage, 3 min)
- Seppu varmam (gluteal region, acupressure, 2 min)
- Nangnananpootu (supra-pubic, light tapping, 2 min)
- Koombu varmam region (calf region, deep pressure, 2 min)

Dietary Counselling

Weekly 30-minute sessions focused on:

- Reduction of refined carbohydrates and processed foods.
- Increased consumption of fruits, vegetables, whole grains, sprouts, and legumes.
- Adequate hydration (2-3 liters water daily).
- Regular meal timing and portion control.
- Elimination of trans fats and fried foods.

Lifestyle Modification

Participants were encouraged to:

- Maintain regular sleep patterns (7-8 hours nightly).
- Engage in daily physical activity (30-45 minutes walking).
- Practice stress management techniques.
- Avoid alcohol and tobacco

Statistical Analysis

Data analysis was performed using SPSS Statistics for Windows (Version 25.0, IBM Corp., and Armonk, NY, USA). Descriptive statistics (mean, standard deviation, frequencies, and percentages) were calculated for all variables. Paired t-tests were used to compare continuous variables pre- and post-intervention. McNemar's test was used for categorical variables (ultrasonography grade). Significance level was set at $\alpha=0.05$ (two-tailed). Effect sizes were calculated using Cohen's d (interpretation: small 0.2-0.5, medium 0.5-0.8, large >0.8).

RESULTS

Participant Characteristics

A total of 30 participants (18 females, 12 males) completed the study with no dropouts. Baseline characteristics are presented in Table 1.

Table 1: Baseline Characteristics of Participants (N=30)

Characteristic	Category	Mean ± SD / n (%)
Age (years)		42.7 ± 8.6
Gender	Male	12 (40.0%)
	Female	18 (60.0%)
Marital Status	Married	24 (80.0%)
	Unmarried	6 (20.0%)
Teaching Experience (years)		15.4 ± 9.2
Institution Type	Government	16 (53.3%)
	Private	14 (46.7%)
NAFLD Grade	Grade I	20 (66.7%)
	Grade II	10 (33.3%)
BMI (kg/m^2)		28.4 ± 4.2
Waist Circumference (cm)		94.3 ± 10.8

ALT (U/L)		54.6 ± 22.3
AST (U/L)		46.8 ± 18.5
PSS-10 Score		24.6 ± 6.8
WHOQOL-BREF	Physical Domain	52.3 ± 12.4
	Psychological Domain	54.8 ± 11.9
	Social Domain	60.2 ± 13.6
	Environmental Domain	58.7 ± 14.2

Abbreviations: NAFLD, Non-Alcoholic Fatty Liver Disease; BMI, Body Mass Index; ALT, Alanine Aminotransferase; AST, Aspartate Aminotransferase; PSS-10, Perceived Stress Scale-10; WHOQOL-BREF, World Health Organization Quality of Life-BREF.

Changes in Liver Function Parameters

Post-intervention analysis revealed statistically significant reductions in both serum ALT and AST.

Table 2: Changes in Liver Function Parameters

Parameter	Pre-Intervention (Mean ± SD)	Post-Intervention (Mean ± SD)	Mean Change	t-value	p-value*	Effect Size (d)
ALT (U/L)	54.6 ± 22.3	34.2 ± 14.8	-20.4	6.82	<0.001	1.10
AST (U/L)	46.8 ± 18.5	30.6 ± 12.1	-16.2	5.94	<0.001	0.96

Abbreviations: ALT, Alanine Aminotransferase; AST, Aspartate Aminotransferase; SD, Standard Deviation.

Statistically significant at $p < 0.05$. Effect size interpretation: small (0.2–0.5), medium (0.5–0.8), large (>0.8).

5.3 Changes in Anthropometric Parameters

Significant improvements were observed in both BMI and waist circumference.

Table 3: Changes in Anthropometric Parameters

Parameter	Pre-Intervention (Mean ± SD)	Post-Intervention (Mean ± SD)	Mean Change	t-value	p-value*	Effect Size (d)
BMI (kg/m^2)	28.4 ± 4.2	26.8 ± 3.9	-1.6	5.42	<0.001	0.89
Waist Circumference (cm)	94.3 ± 10.8	90.1 ± 10.2	-4.2	4.87	<0.001	0.82

Abbreviations: BMI, Body Mass Index; SD, Standard Deviation.

Statistically significant at $p < 0.05$. Effect size interpretation: small (0.2–0.5), medium (0.5–0.8), large (>0.8).

Changes in Perceived Stress

Perceived Stress Scale-10 scores demonstrated a significant reduction following the intervention.

Table 4: Changes in Perceived Stress

Parameter	Pre-Intervention (Mean ± SD)	Post-Intervention (Mean ± SD)	Mean Change	t-value	p-value*	Effect Size (d)
PSS-10 Score	24.6 ± 6.8	16.4 ± 5.2	-8.2	7.24	<0.001	1.32

Abbreviations: PSS-10, Perceived Stress Scale-10; SD, Standard Deviation.

Statistically significant at $p < 0.05$. Effect size interpretation: small (0.2–0.5), medium (0.5–0.8), large (>0.8).

Changes in Quality of Life

Significant improvements were observed across all domains of the WHOQOL-BREF.

Table 5: Changes in Quality of Life (WHOQOL-BREF)

Domain	Pre-Intervention (Mean ± SD)	Post-Intervention (Mean ± SD)	Mean Change	t-value	p-value*	Effect Size (d)
Physical	52.3 ± 12.4	68.6 ± 14.2	+16.3	5.82	<0.001	1.02
Psychological	54.8 ± 11.9	70.2 ± 13.8	+15.4	5.63	<0.001	0.98
Social	60.2 ± 13.6	73.8 ± 15.1	+13.6	4.94	<0.001	0.85
Environmental	58.7 ± 14.2	72.4 ± 15.6	+13.7	4.76	<0.001	0.82

Abbreviations: WHOQOL-BREF, World Health Organization Quality of Life-BREF; SD, Standard Deviation.
Statistically significant at $p < 0.05$. Effect size interpretation: small (0.2–0.5), medium (0.5–0.8), large (>0.8).

5.6 Changes in Ultrasonographic Fatty Liver Grade

Ultrasonographic assessment demonstrated significant improvement in fatty liver grading.

Table 6: Changes in Ultrasonographic Fatty Liver Grade

Grade	Pre-Intervention n (%)	Post-Intervention n (%)	Change
Normal	0 (0.0%)	12 (40.0%)	+12
Grade I	20 (66.7%)	13 (43.3%)	-7
Grade II	10 (33.3%)	5 (16.7%)	-5

Abbreviations: n, number of participants.

Note: Among participants with Grade II NAFLD at baseline (n=10), 6 (60%) showed regression to Grade I or normal. Among Grade I participants (n=20), 12 (60%) showed regression to normal liver status. Overall, 25 out of 30 participants (83.3%) demonstrated improvement in fatty liver grade. McNemar's test revealed significant improvement ($p < 0.001$).

Summary of Findings

The integrated intervention produced significant improvements across all measured outcomes:

Outcome Measure	Percentage Change	Clinical Significance
ALT Reduction	37.4%	Substantial improvement in hepatocyte integrity
AST Reduction	34.6%	Substantial improvement in hepatocyte integrity
BMI Reduction	5.6%	Clinically significant weight loss
Waist Circumference Reduction	4.5%	Reduced visceral adiposity
PSS-10 Reduction	33.3%	Marked reduction in perceived stress
Quality of Life Improvement	26.7–31.2%	Enhanced overall well-being
Radiological Improvement	83.3%	Disease reversal in majority of participants

Abbreviations: ALT, Alanine Aminotransferase; AST, Aspartate Aminotransferase; BMI, Body Mass Index; PSS-10, Perceived Stress Scale-10.

DISCUSSION

Interpretation of Findings

This study demonstrates that an integrated mind-body intervention combining Raja Yoga Meditation, Yoga, Varma therapy, dietary counselling, and lifestyle modification produces significant improvements across multiple domains relevant to NAFLD management among teachers.

Hepatic Function

"The significant reductions in serum ALT and AST levels (37.4% and 34.6%, respectively) indicate substantial improvement in hepatocyte integrity and liver function. These findings align with previous research demonstrating that lifestyle interventions improve liver enzymes in NAFLD patients (Kenneally et al., 2017; Younossi et al., 2016; Younossi et al., 2023). The magnitude of enzyme reduction observed in this study is comparable to or exceeds that reported in pharmacological trials, suggesting that the integrated intervention may offer substantial therapeutic benefit.

The mechanisms underlying hepatic improvement are likely multifactorial. Yoga may enhance hepatic blood flow, improve metabolic regulation, and reduce hepatic inflammation through autonomic modulation and reduced oxidative stress. Dietary modifications directly reduce hepatic fat accumulation and inflammatory burden. Raja Yoga Meditation may reduce cortisol secretion and improve

insulin sensitivity, consequently reducing hepatic fat deposition and hepatocyte damage.

Anthropometric Improvements

"The reductions in BMI (5.6%) and waist circumference (4.5%) are clinically significant, as even modest weight loss (3-5%) has been shown to reduce hepatic steatosis (Kenneally et al., 2017; Cusi, 2016)."The observed improvements likely reflect enhanced energy expenditure through Yoga practice, improved dietary adherence, and reduced stress-related eating behaviors.

The reduction in waist circumference is particularly noteworthy, as visceral adiposity is a key determinant of metabolic dysfunction and NAFLD severity. Decreased central obesity contributes to improved insulin sensitivity, reduced inflammatory mediators, and lower cardiovascular risk.

Stress Reduction

The substantial reduction in perceived stress (33.3%) represents one of the most significant findings of this study. Chronic stress contributes to NAFLD progression through multiple pathways including HPA axis dysregulation, increased cortisol secretion, insulin resistance, and adverse behavioural changes. Raja Yoga Meditation specifically has demonstrated effects on autonomic function and stress regulation.

The stress-reducing effects likely mediated improvements in metabolic parameters through reduced sympathetic activation, improved emotional regulation, and decreased stress-induced hyperphagia. Additionally, reduced stress may have enhanced participants' capacity for self-regulation, supporting adherence to dietary and lifestyle recommendations.

Quality of Life

Significant improvements across all WHOQOL-BREF domains indicate that the intervention enhanced overall well-being beyond metabolic outcomes. "This is consistent with research demonstrating that mind-body interventions improve health-related quality of life in various populations (Cramer et al., 2016; Pascoe & Bauer, 2015)." Improved physical health, reduced psychological distress, enhanced social functioning, and better environmental satisfaction collectively contributed to improved quality of life.

Ultrasonographic Regression

The radiological improvements, with 83.3% of participants showing reduced fatty liver grade and 40% achieving normal liver status, represent clinically meaningful disease reversal. These findings suggest that the integrated intervention may address the underlying pathophysiological mechanisms of NAFLD rather than merely ameliorating symptoms.

Mechanisms of Action

The therapeutic benefits observed likely arise from synergistic effects of multiple intervention components:

Yoga: Increased energy expenditure and physical activity; improved insulin sensitivity and glucose metabolism; enhanced autonomic balance and reduced sympathetic overactivity; improved circulation and hepatic blood flow; reduced inflammation and oxidative stress.

Raja Yoga Meditation: Reduced stress through downregulation of HPA axis; decreased cortisol and catecholamine secretion; improved emotional regulation and psychological resilience; enhanced parasympathetic activity; improved self-regulation and health behaviours.

Varma Therapy: Potential regulation of autonomic and circulatory function; stimulation of specific neural and energetic pathways; enhanced relaxation and stress reduction; possible modulation of organ-specific function.

Dietary Counselling: Reduced caloric and carbohydrate intake; decreased saturated fat and processed food consumption; increased antioxidant and fiber intake; improved glycemic control and insulin sensitivity.

Lifestyle Modification: Enhanced physical activity and energy expenditure; improved sleep quality and circadian regulation; reduced sedentary behavior; enhanced self-care and health awareness.

Comparison with Previous Studies

"The findings are consistent with previous research demonstrating that yoga and meditation improve metabolic profiles, reduce stress, and enhance quality of life (Pascoe & Bauer, 2015; Goyal et al., 2014; Black & Slavich, 2016; Cramer et al., 2016)." However, few studies have specifically evaluated integrated mind-body interventions in NAFLD, and fewer have focused on teachers as a vulnerable occupational group.

Compared to pharmacological interventions for NAFLD, the integrated mind-body approach offers advantages including no adverse effects or drug-related toxicity, benefits across multiple physiological and psychological domains, long-term sustainability and self-management potential, and cost-effectiveness and accessibility.

Clinical and Public Health Implications

The findings have important implications:

Occupational Health: Workplace-based integrative wellness programs may effectively address NAFLD burden among teachers. Schools and educational institutions could consider implementing similar programs to improve teacher health and well-being.

Healthcare Delivery: Mind-body interventions can complement conventional management, potentially reducing medication requirements and healthcare utilization. Integrative approaches may be particularly valuable for patients who do not respond to or cannot tolerate pharmacological interventions.

Health Promotion: Integrated approaches addressing both physical and psychological determinants may improve adherence and outcomes. The multimodal nature of the intervention may be more engaging and sustainable than single-component approaches.

Resource Allocation: Cost-effective interventions may reduce the economic burden of NAFLD. The low-cost nature of mind-body therapies makes them accessible for implementation in resource-limited settings.

CONCLUSION

Summary of Findings

This study provides evidence that an integrated intervention comprising Raja Yoga Meditation, Yoga, Varma therapy, dietary counselling, and lifestyle modification produces significant and clinically meaningful improvements in liver function, anthropometric parameters, perceived stress, quality of life, and fatty liver grade among secondary school teachers with NAFLD. The findings support the alternative hypothesis and reject the null hypothesis.

Significance and Contributions

The study contributes to the growing body of evidence supporting integrative mind-body therapies for metabolic disorders. Key contributions include:

- Demonstration of effective NAFLD management in a vulnerable occupational group.
- Evidence for the effectiveness of an integrated rather than single-modality intervention.
- Support for incorporating traditional mind-body therapies in contemporary healthcare.
- Documentation of both physiological and psychological benefits.
- Identification of a potentially scalable intervention for workplace settings.

Implications

The findings highlight the importance of incorporating integrative mind-body therapies into comprehensive healthcare programs for NAFLD. Occupational health interventions targeting teachers should consider the combined implementation of Yoga, Raja Yoga Meditation, Varma therapy, and lifestyle modification to address the multifactorial etiology of NAFLD. Policymakers and healthcare administrators should consider supporting the integration of such approaches into public health programs.

Limitations and Future Research

Limitations

- Study Design: The quasi-experimental design (one-group pre-post) without a control group limits causal attribution. Future research employing randomized controlled trials with active comparators is warranted.
- Sample Size: The small sample size (n=30) limits statistical power and generalizability. Larger studies with adequate power are needed to confirm findings and enable subgroup analyses.
- Sample Characteristics: Participants were recruited from a single geographical region, limiting generalizability to other populations and settings. Multicenter studies are needed.
- Disease Severity: The study focused on Grade I and Grade II NAFLD only; findings may not apply to more severe disease (Grade III, NASH, fibrosis).
- Intervention Duration: The 8-week intervention may not be sufficient to assess long-term effects and sustainability of improvements. Extended follow-up periods (6-12 months) are recommended.
- Confounding Variables: Lifestyle factors outside the intervention could not be fully controlled, potentially influencing outcomes.
- Adherence Variability: Participant adherence to the intervention may have varied, potentially affecting outcomes.
- Therapy Specificity: The study evaluated an integrated intervention and not the isolated effects of each therapy, limiting the ability to attribute benefits to specific components.
- Self-Report Measures: Reliance on self-report questionnaires (PSS-10, WHOQOL-BREF) introduces potential for response bias.
- Blinding: Participants and intervention providers were not blinded to the intervention.

Suggestions for Future Research

Methodological Enhancements

- Conduct randomized controlled trials with larger sample sizes and active comparators.
- Use multicenter designs involving participants from different regions.
- Include wider age range and disease severity.
- Conduct long-term follow-up studies (6-12 months).

Mechanistic Investigations

- Examine inflammatory biomarkers (TNF-α, IL-6), insulin resistance markers (HOMA-IR), oxidative stress markers, and adipokines.
- Measure cortisol and catecholamine levels to evaluate stress-related mechanisms.
- Assess heart rate variability and autonomic balance changes.

- Conduct metabolomic profiling to investigate changes in lipid and glucose metabolism.

Comparative Studies

- Compare individual and combined effects of Raja Yoga Meditation, Yoga, and Varma therapy
- Evaluate optimal intervention duration, frequency, and intensity
- Develop standardized protocols for Varma therapy

Implementation Research

- Develop and evaluate workplace-based integrative wellness programs for teachers.
- Investigate barriers and facilitators to implementation in educational settings.
- Conduct cost-effectiveness analyses to inform healthcare policy decisions.
- Assess impact on occupational performance and productivity.

Advanced Diagnostics

- Utilize quantitative assessments such as FibroScan, MRI-PDFF, or transient elastography for objective hepatic fat quantification.
- Examine changes in liver stiffness as a marker of fibrosis progression.

REFERENCES

1. Younossi, Z. M., Koenig, A. B., Abdelatif, D., et al. (2016). Global epidemiology of non-alcoholic fatty liver disease- Meta-analytic assessment of prevalence, incidence, and outcomes. Hepatology, 64(1), 73-84.
2. Cusi, K. (2016). Role of insulin resistance in non-alcoholic fatty liver disease. Journal of Clinical and Endocrinology, 101(3), 1-10.
3. Friedman, S. L., Neuschwander-Tetri, B. A., Rinella, M., & Sanyal, A. J. (2018). Mechanisms of NAFLD development and therapeutic strategies. Nature Medicine, 24(7), 908-922.
4. Younossi, Z. M., et al. (2023). Global burden of fatty liver disease and the need for non-pharmacological interventions. Hepatology.
5. Kenneally, S., Sier, J. H., & Moore, J. B. (2017). Efficacy of dietary and physical activity intervention in non-alcoholic fatty liver disease: A systematic review. BMJ Open Gastroenterology, 4(1), e000139.
6. Kyriacou, C. (2001). Teacher stress: Directions for future research. Educational Review, 53(1), 27-35.
7. Jennings, P. A., & Greenberg, M. T. (2009). The prosocial classroom: Teacher social and emotional competence in relation to student and classroom outcomes. Review of Educational Research, 79(1), 491-525.
8. Roeser, R. W., Skinner, E., Beers, J., & Jennings, P. A. (2013). Mindfulness training and teachers' professional development. Mindfulness, 4(2), 89-102.
9. Pascoe, M. C., & Bauer, I. E. (2015). A systematic review of randomised control trials on the effects of yoga on stress measures and mood. Journal of Psychiatric Research, 68, 270-282.
10. Goyal, M., Singh, S., Sibinga, E. M. S., et al. (2014). Meditation programs for psychological stress and well-being: A systematic review and meta-analysis. JAMA, 312(20), 2131-2138.
11. Black, D. S., & Slavich, G. M. (2016). Mindfulness meditation and the immune system: A systematic review of randomized controlled trials. Annals of the New York Academy of Sciences, 1373(1), 13-24.
12. Sharma, V. K., Das, S., Mondal, S., et al. (2015). Effect of Rajyoga meditation on autonomic functions. International Journal of Yoga, 8(1), 14-18.
13. Eslam, M., Sanyal, A. J., & George, J. (2020). MAFLD: A consensus-driven proposed nomenclature. Journal of Hepatology, 73(1), 202-209.
14. Cramer, H., Lauche, R., Haller, H., & Dobos, G. (2016). Yoga for improving health-related quality of life, mental health and cancer-related symptoms in women diagnosed with breast cancer. Cochrane Database of Systematic Reviews, (1), CD010802.
15. Younossi, Z. M., et al. (2023). Global burden of fatty liver disease and the need for non-pharmacological interventions. Hepatology.
16. Eslam, M., Sanyal, A. J., & George, J. (2020). MAFLD: A consensus-driven proposed nomenclature.

Cite this article as:
S. Venkatesan, D R Rajendran. Integrative Mind-Body Therapies for Non-Alcoholic Fatty Liver Disease: A Pre-Post Interventional Study Among Secondary School Teachers. International Journal of Ayurveda and Pharma Research. 2026;14(8):51-59.
https://doi.org/10.47070/ijapr.v14i8.4287
Source of support: Nil, Conflict of interest: None Declared

***Address for correspondence**
Dr. S. Venkatesan
Research Scholar,
Manipur International University,
Manipur.
Email: venkatesansudalai@gmail.com

Disclaimer: IJAPR is solely owned by Mahadev Publications - dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IJAPR cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of IJAPR editor or editorial board members.